

# NN scattering in higher derivative formulation of baryon chiral perturbation theory

D. Djukanovic,<sup>1</sup> J. Gegelia,<sup>1,2</sup> S. Scherer,<sup>1</sup> and M. R. Schindler<sup>1</sup>

<sup>1</sup>*Institut für Kernphysik, Johannes Gutenberg-Universität, D-55099 Mainz, Germany*

<sup>2</sup>*Institute of High Energy Physics and Informatization,  
Tbilisi State University, 0186 Tbilisi, Georgia*

(Dated: Today)

## Abstract

We consider a new approach to the nucleon-nucleon scattering problem in the framework of the higher-derivative formulation of baryon chiral perturbation theory. Starting with a Lorentz-invariant form of the effective Lagrangian we work out a new symmetry-preserving framework where the leading-order amplitude is calculated by solving renormalizable equations and corrections are taken into account perturbatively. Analogously to the KSW approach, the (leading) renormalization scale dependence to any finite order is absorbed in the redefinition of a finite number of parameters of the effective potential at given order. On the other hand, analogously to Weinberg's power counting, the one-pion-exchange potential is of leading order and is treated non-perturbatively.

PACS numbers: 11.10.Gh,12.39.Fe,13.75.Cs

## I. INTRODUCTION

The few-nucleon sector of baryon chiral perturbation theory was first considered in the seminal papers by Weinberg [1, 2] which triggered an intensive research activity starting with the pioneering work of Ref. [3]. In this approach for processes involving  $N > 1$  nucleons the power counting is applied to the effective potential, which is defined as the sum of all  $N$ -nucleon-irreducible diagrams. The scattering amplitudes are obtained by solving the Lippmann-Schwinger (LS) or Schrödinger equation. For recent reviews and references see e.g. Refs. [4, 5].

The implementation of Weinberg's program encountered problems. The NN potential of the effective field theory (EFT) is non-renormalizable in the traditional sense already at leading order. The iteration of the potential in the LS equation generates divergent terms with structures which are not included in the original potential. Therefore, the renormalization of the solution to the LS equation requires the contributions of an infinite number of higher-order counter-terms. Here, the infinite number refers to both the loop and the chiral expansions. The freedom of choosing the finite parts of these counter-terms is compensated by the running of the corresponding renormalized couplings. It has been argued that the coefficients of the divergent parts of the counter-terms contributing in low-order calculations would set the scale of the corresponding renormalized couplings. As a consequence, even if these couplings were natural at some value of the renormalization scale, they would become unnaturally large for slightly different values of this parameter. This problem (also analyzed non-perturbatively), in different variations, has been addressed as the inconsistency of Weinberg's approach, and an alternative power counting scheme, known as the KSW approach, has been suggested [6, 7, 8, 9]. In the KSW scheme the troublesome one-pion-exchange (OPE) potential is shifted from leading order (LO) to next-to-leading order (NLO). The leading-order equation becomes renormalizable, i.e. all divergences can be absorbed in the redefinition of the parameters of the LO potential, perturbatively as well as non-perturbatively. Moreover, the LO equation is exactly solvable and dimensional regularization can be applied. Corrections are treated perturbatively in the KSW approach. This guarantees that *all* divergences are absorbed in parameters contributing to the given order. As a consequence, the KSW approach is free of "consistency problems." Unfortunately this scheme suffers from convergence problems. As the OPE potential is shifted to NLO, it is treated perturbatively. It has been argued that the perturbative treatment of pions in NN scattering poses problems of convergence [10, 11, 12]. The results of Ref. [13] made it explicit that perturbative series do not converge in the KSW approach, at least in the triplet channel.

As a next step, in Ref. [14] a non-perturbative investigation of the renormalization problem has been carried out and an expansion of nuclear forces about the chiral limit has been suggested. It has been argued that this expansion is formally consistent and is equivalent to the KSW power counting in the  $^1S_0$  channel and to Weinberg's power counting in the  $^3S_1 - ^3D_1$  coupled channels. Higher partial waves have been investigated in Ref. [15] with the result that the perturbative treatment of the OPE potential is not sufficient for a finite number of partial waves, e.g.  $^3P_0$  and  $^3P_2 - ^3F_2$ . On the other hand, the iterated OPE potential produces a cutoff dependence in all waves where the tensor force is attractive. This requires additional counter-terms in each of these partial waves, clearly upsetting the power counting. The authors of Ref. [15] conjecture that the "mixture of perturbative treatment of higher partial waves, resummation of lower partial waves, and promotion of a finite

number of counter-terms is the most consistent approach” to the chiral perturbation theory for nuclear systems while sub-leading interactions are treated perturbatively.

There exists an alternative point of view that the ”consistency problem” of Weinberg’s approach is irrelevant and one can perform the calculations in cutoff effective field theory (EFT) by suitably choosing the cutoff parameter. According to this point of view, taking large values of the cutoff parameter without including the contributions of all (perturbatively) relevant counter-terms is not a legitimate procedure. Instead one has to choose the cutoff parameter of the order of the relevant large scale, presumably the mass of the  $\rho$  meson [16, 17, 18, 19, 20, 21, 22, 23].

In the present work we develop a new approach which combines aspects of both points of view outlined above. In particular, in our new framework the leading-order amplitude is calculated by solving integral equations and corrections are taken into account perturbatively. Analogously to the KSW approach, the (leading) renormalization scale dependence to any given finite order is absorbed in the redefinition of a finite number of parameters of the effective potential at the given order. On the other hand, analogously to Weinberg’s scheme, the OPE potential is of leading order and is treated non-perturbatively.

In the two-nucleon sector the path integral for the generating functional of the Green’s functions is not Gaussian and therefore the standard heavy-baryon reduction can not be carried out. While the non-relativistic effective Lagrangian can be fixed by matching to the (explicitly) Lorentz-invariant Lagrangian, we prefer to work in terms of the latter. We start with the higher-derivative formulation of the manifestly Lorentz-invariant Lagrangian of Ref. [24] and perform the non-relativistic reduction at the level of Feynman diagrams.

Our work is organized as follows: In Sec. II the effective Lagrangian in its manifestly Lorentz-invariant form and field redefinitions leading to the higher-derivative formulation are considered. Section III addresses the issues of renormalization and power counting. In Sec. IV the NN scattering equations are derived and analyzed. A summary is given in Sec. V.

## II. THE EFFECTIVE LAGRANGIAN

In this section we consider the effective Lagrangian needed for the purposes of this work. The standard effective Lagrangian is organized in a derivative and quark-mass expansion and consists of the sum of the purely mesonic, the  $\pi N$ ,  $NN$ ,  $\dots$  parts,

$$\mathcal{L}_{\text{eff}} = \mathcal{L}_\pi + \mathcal{L}_{\pi N} + \mathcal{L}_{NN} + \dots \quad (1)$$

The lowest-order mesonic Lagrangian reads [25]

$$\mathcal{L}_2 = \frac{F^2}{4} \text{Tr} \left[ D_\mu U (D^\mu U)^\dagger \right] + \frac{F^2}{4} \text{Tr} (\chi U^\dagger + U \chi^\dagger), \quad (2)$$

where  $U$  is a unimodular unitary ( $2 \times 2$ ) matrix containing the pion fields. The covariant derivative is defined as

$$D_\mu U = \partial_\mu U - i r_\mu U + i U l_\mu,$$

and

$$r_\mu = v_\mu + a_\mu, \quad l_\mu = v_\mu - a_\mu, \quad \chi = 2B(s + ip).$$

Here,  $v_\mu$ ,  $a_\mu$ ,  $s$ , and  $p$  are external vector, axial-vector, scalar, and pseudo-scalar sources, respectively. In Eq. (2),  $F$  denotes the pion-decay constant in the chiral limit:  $F_\pi = F[1 + \mathcal{O}(\hat{m})] = 92.4$  MeV. We consider the isospin-symmetric limit  $m_u = m_d = \hat{m}$ , and the lowest-order expression for the squared pion mass is  $M^2 = 2B\hat{m}$ , where  $B$  is related to the quark condensate  $\langle \bar{q}q \rangle_0$  in the chiral limit [25].

The nucleon field is described with two four-component Dirac fields,

$$\Psi = \begin{pmatrix} p \\ n \end{pmatrix},$$

and the lowest-order Lagrangian of the one-nucleon sector reads [26]

$$\mathcal{L}_{\pi N}^{(1)} = \bar{\Psi} \left( i\gamma_\mu D^\mu - m + \frac{1}{2} \overset{\circ}{g}_A \gamma_\mu \gamma_5 u^\mu \right) \Psi, \quad (3)$$

where  $D_\mu \Psi = (\partial_\mu + \Gamma_\mu - i v_\mu^{(s)}) \Psi$  denotes the covariant derivative with the iso-scalar vector source  $v_\mu^{(s)}$ , and the quantities  $u$ ,  $u_\mu$ ,  $\Gamma_\mu$ , and  $\chi_\pm$  (and their derivatives) are defined as

$$u^2 = U, \quad u_\mu = iu^\dagger D_\mu U u^\dagger, \quad \Gamma_\mu = \frac{1}{2} [u^\dagger \partial_\mu u + u \partial_\mu u^\dagger - i(u^\dagger r_\mu u + u l_\mu u^\dagger)],$$

$$\chi_\pm = u^\dagger \chi u^\dagger \pm u \chi^\dagger u.$$

In Eq. (3),  $m$  and  $\overset{\circ}{g}_A$  stand for the chiral limit of the physical nucleon mass and the axial-vector coupling constant, respectively.

The most general chirally-invariant leading-order NN Lagrangian reads (for the power counting see below)

$$\begin{aligned} \mathcal{L}_{NN} = \sum_{a=0}^3 & \left[ C_S^a \bar{\Psi} \tau^a \Psi \bar{\Psi} \tau^a \Psi + C_A^a \bar{\Psi} \tau^a \gamma_5 \Psi \bar{\Psi} \tau^a \gamma_5 \Psi + C_V^a \bar{\Psi} \tau^a \gamma_\mu \Psi \bar{\Psi} \tau^a \gamma^\mu \Psi \right. \\ & \left. + C_{AV}^a \bar{\Psi} \tau^a \gamma_5 \gamma_\mu \Psi \bar{\Psi} \tau^a \gamma_5 \gamma^\mu \Psi + C_T^a \bar{\Psi} \tau^a \sigma_{\mu\nu} \Psi \bar{\Psi} \tau^a \sigma^{\mu\nu} \Psi \right], \end{aligned} \quad (4)$$

where  $\tau^0$  is the unit (isospin) matrix and  $\tau^i$  ( $i = 1, 2, 3$ ) are (isospin) Pauli matrices and  $C_I^1 = C_I^2 = C_I^3$  for all  $I$ . Note that the structure  $\bar{\Psi} \gamma_5 \Psi$  is assigned leading order.

The most general effective Lagrangian includes *all* terms which are consistent with the symmetries of the underlying fundamental theory, i.e. QCD. The above given "canonical" form of the effective Lagrangian is obtained from the most general form by using suitable field redefinitions [27]. This is done for convenience. Any two forms of the effective Lagrangian obtained from each other by using field redefinitions are completely equivalent when physical quantities are considered. Note that the coupling constants corresponding to the same operator structures in two forms of the effective Lagrangian are in general not equal [28]. Switching from one formulation to the other, one needs to take into account the new contributions (in general in all operator structures) generated by field transformations. There is an infinite number of possible field transformations and correspondingly an infinite number of forms of the effective Lagrangian. While all these forms lead to the same expressions for physical quantities (with certain care), in practical (perturbative) calculations one form of the effective Lagrangian could be considerably more convenient than the other.

Starting with the above given "canonical" effective Lagrangian one encounters ultraviolet divergences in the equations for the few-nucleon scattering amplitudes. This problem can be handled by considering a nucleon propagator with improved ultraviolet behavior [24]. To that end we employ a "non-canonical" form of the effective Lagrangian generated by using suitable field transformations. To obtain the modified nucleon propagator we perform the following field transformations

$$\begin{aligned}\Psi &\rightarrow \Phi \left( L^2, -\vec{D}^2 \right) \Psi, \\ \bar{\Psi} &\rightarrow \bar{\Psi} \Phi \left( L^2, -\overleftarrow{D}^2 \right),\end{aligned}\tag{5}$$

where  $\bar{\Psi} \overleftarrow{D}^2 = \left( \vec{D}^2 \Psi \right)^\dagger \gamma^0$  and obtain

$$\mathcal{L}_{\pi N}^{(1)'} = \bar{\Psi} \Phi \left( L^2, -\overleftarrow{D}^2 \right) \left( i\gamma_\mu D^\mu - m + \frac{1}{2} \overset{\circ}{g}_A \gamma_\mu \gamma_5 u^\mu \right) \Phi \left( L^2, -\vec{D}^2 \right) \Psi.\tag{6}$$

In Eq. (6)  $L$  is a free parameter,  $\vec{D}$  stands for the spacial components of the covariant derivative and  $\Phi$  is some appropriately chosen real function. Possible choices of  $\Phi$  are, e.g.,

$$\Phi \left( L^2, X \right) = \left( 1 + \frac{1}{L^2} X \right)^{N_\Psi} \text{ or } \Phi \left( L^2, X \right) = \exp \left( \frac{1}{L^2} X \right)^{N_\Psi}.\tag{7}$$

The Feynman propagator corresponding to the Lagrangian of Eq. (6) reads

$$i S_F(p) = \frac{i}{\not{p} - m + i\epsilon} F(L^2, \vec{p}^2),\tag{8}$$

with

$$F(L^2, \vec{p}^2) = \frac{L^{4N_\Psi}}{(L^2 + \vec{p}^2)^{2N_\Psi}} \text{ and } F(L^2, \vec{p}^2) = \exp \left\{ -2 \left( \frac{\vec{p}^2}{L^2} \right)^{N_\Psi} \right\},\tag{9}$$

respectively.

Depending on the order of the performed calculations we choose the modified propagator such that all loop integrations over the momenta of the intermediate purely nucleonic states are rendered finite. We apply dimensional regularization to the resulting effective theory, expand the regularized diagrams in powers of  $n-4$  and subtract  $1/(n-4)$  pole-terms, where  $n$  is the space-time dimension. There is a finite number of pole-terms to any given (finite) order in the chiral expansion of physical quantities in the vacuum and one-nucleon sectors, and the effective potentials in the two-nucleon sector. No further divergences occur (for finite parameters  $L$ ) neither in the vacuum and one-nucleon sector nor in the equations of the two-nucleon sector. Therefore, we can take  $n=4$  in the equations of the two-nucleon sector after renormalizing the effective potential and the corrections to the two-nucleon propagator. Note that the field redefinition of Eq. (5) introduces additional interaction terms in the effective Lagrangian. Their contributions are important for preserving the symmetries of the theory. In practice it is convenient to choose the exponential form of the field redefinition with sufficiently large  $N_\Psi$  [5] where the additional interaction terms start contributing at orders higher than the order of accuracy of the performed calculations.

The complete dependence of the physical quantities on the parameter  $L$  can be absorbed in the redefinition of (an infinite number of) coupling constants. Therefore different choices of

the numerical value of this parameter correspond to different renormalization schemes. Note that the applied regularization scheme is the dimensional regularization. The parameter  $L$  should not be confused with the cutoff parameter and it does not have to be taken to infinity.

### III. POWER COUNTING

It is understood that the power counting specified below is satisfied by renormalized diagrams, provided the renormalization points are chosen appropriately. In particular, following Ref. [2] one should choose the renormalization points of the order of small external momenta.

#### A. Effective potential

Interaction terms, generated by the one-nucleon sector of the effective Lagrangian [29] are assigned the standard orders of BChPT [30]. In the two-nucleon sector each derivative acting on the pion field as well as each factor of the pion mass is assigned order  $q^1$ , where  $q$  denotes a small quantity like the pion mass or a nucleon three momentum. Contracted derivatives acting on nucleon fields lead to scalar products of large nucleon momenta, e.g.  $p_1 \cdot p_2$ . We can rewrite

$$2p_1 \cdot p_2 = (p_1^2 - m^2) + (p_2^2 - m^2) - (p_1 - p_2)^2 + 2m^2. \quad (10)$$

As in the one-nucleon sector,  $(p_i^2 - m^2)$  is counted as  $q^1$ , while the difference  $(p_1 - p_2)^2$  is of order  $q^2$ . We arrange the Lagrangian such that terms involving derivatives are accompanied by interaction terms where each pair of contracted derivatives is replaced with a factor of " $+m^2$ " or " $-m^2$ ". Therefore each pair of contracted derivatives acting on nucleon fields is assigned order  $q^1$  (we choose the renormalization scheme such that this power counting also holds for loop diagrams.) For example, the term  $\bar{\Psi} D_\mu \Psi \bar{\Psi} D^\mu \Psi$  is rewritten as

$$(\bar{\Psi} D_\mu \Psi \bar{\Psi} D^\mu \Psi + m^2 \bar{\Psi} \Psi \bar{\Psi} \Psi) - m^2 \bar{\Psi} \Psi \bar{\Psi} \Psi, \quad (11)$$

where the first and second terms are part of the NLO and LO NN Lagrangian, respectively.

We assign to each diagram contributing to the effective potential a chiral order  $D$  determined as follows. Vertices from the  $k$ -th order Lagrangian count as order  $q^k$ , a loop integration in  $n$  dimensions as  $q^n$ , each pion propagator as  $q^{-2}$  and each nucleon propagator as  $q^{-1}$ , respectively. Note that the terms of the  $k$ -th order NN Lagrangian can only generate contributions to effective potential of order  $k$  and higher. To any finite order there is only a finite number of terms in the two-nucleon sector of the effective Lagrangian. Correspondingly the effective potential contains a finite number of terms to any finite order. We drop nucleon loops, including their contributions in the redefinition of the coupling constants of the effective Lagrangian.

#### B. Two-nucleon propagator

We rewrite the fermion propagator as

$$S_F(p) = \frac{\not{p} + m}{p^2 - m^2 + i\epsilon} F(L^2, \vec{p}^2)$$

$$\begin{aligned}
&= \frac{m P_+ F(L^2, \vec{p}^2)}{\sqrt{\vec{p}^2 + m^2} (p_0 - \sqrt{\vec{p}^2 + m^2} + i\epsilon)} - \frac{m P_+ F(L^2, \vec{p}^2)}{\sqrt{\vec{p}^2 + m^2} (p_0 + \sqrt{\vec{p}^2 + m^2} - i\epsilon)} \\
&\quad + \frac{(\not{p} - m \not{\psi}) F(L^2, \vec{p}^2)}{2 \sqrt{\vec{p}^2 + m^2} (p_0 - \sqrt{\vec{p}^2 + m^2} + i\epsilon)} - \frac{(\not{p} - m \not{\psi}) F(L^2, \vec{p}^2)}{2 \sqrt{\vec{p}^2 + m^2} (p_0 + \sqrt{\vec{p}^2 + m^2} - i\epsilon)} \\
&= S_F^{(1)}(p) + S_F^{(2)}(p) + S_F^{(3)}(p) + S_F^{(4)}(p), \tag{12}
\end{aligned}$$

where the projector  $P_+$  is defined as

$$P_+ = \frac{1 + \not{v}}{2}, \tag{13}$$

with  $v = (1, 0, 0, 0)$ . The poles in  $S_F^{(1)}$  and  $S_F^{(3)}$  correspond to particle states, and those in  $S_F^{(2)}$  and  $S_F^{(4)}$  to anti-particles, respectively. We assign order  $q^{-1}$  to  $S_F^{(1)}$ , order  $q^0$  to the second and third terms  $S_F^{(2)}$  and  $S_F^{(3)}$ , and order  $q^1$  to  $S_F^{(4)}$ .

The undressed two-nucleon propagator is defined as the product of two undressed one-nucleon propagators. Inserting the expansion of Eq. (12) we obtain a sum of terms of different orders. For nucleon-nucleon scattering the products of  $S_F^{(1)}$  and  $S_F^{(3)}$  are enhanced in purely two-nucleon intermediate states [1, 2]. Therefore, we assign the following orders to the various terms in the undressed two-nucleon propagator:

$$\begin{aligned}
S_F^{(1)} S_F^{(1)} &\sim q^{-4}, & S_F^{(1)} S_F^{(3)} &\sim q^{-3}, \\
S_F^{(2)} S_F^{(2)} &\sim q^0, & S_F^{(2)} S_F^{(4)} &\sim q^1, \\
S_F^{(3)} S_F^{(3)} &\sim q^{-2}, & S_F^{(4)} S_F^{(4)} &\sim q^2, \\
S_F^{(1)} S_F^{(2)} &\sim q^{-1}, & S_F^{(1)} S_F^{(4)} &\sim q^0, \\
S_F^{(2)} S_F^{(3)} &\sim q^0, & S_F^{(3)} S_F^{(4)} &\sim q^1.
\end{aligned} \tag{14}$$

The first term in Eq. (14) defines the leading-order two-nucleon propagator  $G_0$ ,

$$G_0 = -i S_F^{(1)} S_F^{(1)}. \tag{15}$$

The last four terms in Eq. (14) possess poles on the same side of the real axis in the complex plane of the zeroth component of the integration momenta. Therefore, if the effective potential does not contain singularities in the complex energy plane, they give vanishing contributions.

#### IV. NN SCATTERING EQUATION

Defining the two-nucleon-irreducible diagrams as the effective potential  $V$  the equation for the amplitude (amputated Green's function) reads

$$\begin{aligned}
T_{\lambda\sigma,\mu\nu}(p_3, p_4, p_1, p_2) &= V_{\lambda\sigma,\mu\nu}(p_3, p_4, p_1, p_2) \\
&\quad - i \int \frac{d^4 k}{(2\pi)^4} V_{\lambda\sigma,\alpha\gamma}(p_3, p_4, p_3 - k, p_4 + k) \\
&\quad \times S_{F,\alpha\beta}(p_3 - k) S_{F,\gamma\delta}(p_4 + k) T_{\beta\delta,\mu\nu}(p_3 - k, p_4 + k, p_1, p_2), \tag{16}
\end{aligned}$$

where the Greek indices labeling the components of the Dirac fermion field run from 1 to 4. The corresponding physical amplitude is obtained as

$$\sqrt{Z_\psi(p_3)} \sqrt{Z_\psi(p_4)} \bar{u}_\lambda(p_3) \bar{u}_\sigma(p_4) T_{\lambda\sigma,\mu\nu}(p_3, p_4, p_1, p_2) u_\mu(p_1) u_\nu(p_2) \sqrt{Z_\psi(p_1)} \sqrt{Z_\psi(p_2)}, \quad (17)$$

where  $Z_\psi$  is the residue of the nucleon propagator. The residue corresponding to the propagator of Eqs. (8) and (12) is

$$Z_\psi(p) = F(L^2, \vec{p}^2). \quad (18)$$

For convenience in the following derivations we write Eq. (16) symbolically as

$$T_{\lambda\sigma,\mu\nu} = V_{\lambda\sigma,\mu\nu} + V_{\lambda\sigma,\alpha\gamma} G_{\alpha\beta,\gamma\delta} T_{\beta\delta,\mu\nu}, \quad (19)$$

or

$$T = V + V G T. \quad (20)$$

Next, we expand  $T$ ,  $V$ , and  $G$  in small parameters,

$$G = G_0 + G_1 + G_2 + \dots, \quad (21)$$

$$V = V_0 + V_1 + V_2 + \dots, \quad (22)$$

$$T = T_0 + T_1 + T_2 + \dots, \quad (23)$$

where the quantities  $X_i$  ( $X = T, G, V$ ) are of order  $\mathcal{O}(q^i)$ . We then substitute the expansions in Eq. (20), and solve  $T$  order by order. At leading order we obtain

$$T_0 = V_0 + V_0 G_0 T_0. \quad (24)$$

To determine the physical amplitude order by order we expand the solutions to the Dirac equation in small quantities

$$\begin{aligned} u &= u_0 + u_1 + u_2 + \dots = P_+ u(p) + \frac{\not{p} - p \cdot v}{2m} P_+ u(p) + \dots, \\ \bar{u} &= \bar{u}_0 + \bar{u}_1 + \bar{u}_2 + \dots = \bar{u}(p) P_+ + \bar{u}(p) P_+ \frac{\not{p} - p \cdot v}{2m} + \dots. \end{aligned} \quad (25)$$

Substituting Eqs. (23) and (25) in Eq. (17) we obtain the expansion of the physical amplitude as

$$\begin{aligned} \sqrt{Z_\psi} \sqrt{Z_\psi} \bar{u} \bar{u} T u u \sqrt{Z_\psi} \sqrt{Z_\psi} &= \sqrt{Z_\psi} \sqrt{Z_\psi} [\bar{u}_0 \bar{u}_0 T_0 u_0 u_0 + \bar{u}_1 \bar{u}_0 T_0 u_0 u_0 + \bar{u}_0 \bar{u}_1 T_0 u_0 u_0 \\ &+ \bar{u}_0 \bar{u}_0 T_1 u_0 u_0 + \bar{u}_0 \bar{u}_0 T_0 u_1 u_0 + \bar{u}_0 \bar{u}_0 T_0 u_0 u_1 + \dots] \sqrt{Z_\psi} \sqrt{Z_\psi}. \end{aligned} \quad (26)$$

Taking into account Eq. (25) we see that to calculate the leading-order on-mass-shell amplitude we need the quantity  $\tilde{T}_0 = P_+ P_+ T_0 P_+ P_+$ . Applying the projectors  $P_+$  to Eq. (24) and using (see Eq. (15))

$$G_0 = G_0 P_+ P_+ = P_+ P_+ G_0 = P_+ P_+ G_0 P_+ P_+ \quad (27)$$

we obtain

$$P_+ P_+ T_0 P_+ P_+ = P_+ P_+ V_0 P_+ P_+ + P_+ P_+ V_0 P_+ P_+ G_0 P_+ P_+ T_0 P_+ P_+, \quad (28)$$

which is an integral equation for the projected amplitude  $\tilde{T}_0$ ,

$$\tilde{T}_0 = \tilde{V}_0 + \tilde{V}_0 G_0 \tilde{T}_0, \quad (29)$$

where  $\tilde{V}_0 = P_+ P_+ V_0 P_+ P_+$ .

To calculate  $T_0$  from the solution to Eq. (29) we rewrite Eq. (24) as

$$T_0 = V_0 + V_0 G_0 V_0 + V_0 G_0 T_0 G_0 V_0, \quad (30)$$

which, using Eq. (27), takes the form

$$T_0 = V_0 + V_0 G_0 V_0 + V_0 G_0 \tilde{T}_0 G_0 V_0. \quad (31)$$

The leading-order amplitude  $T_0$  calculated from Eq. (31) can then be used in calculations of the next-to-leading-order amplitude  $T_1$ . The equation for  $T_1$

$$T_1 = V_1 + V_1 G_0 T_0 + V_0 G_1 T_0 + V_0 G_0 T_1 \quad (32)$$

can be solved exactly. The solution reads

$$T_1 = V_1 + T_0 G_0 V_1 + V_1 G_0 T_0 + T_0 G_0 V_1 G_0 T_0 + T_0 G_1 T_0. \quad (33)$$

Analogously to the above case the leading-order amplitude  $T_0$  and the next-to-leading-order amplitude  $T_1$  can be used to calculate the next-to-next-to-leading-order amplitude  $T_2$  etc.

### A. Leading-order equation

In the following, we will consider the scattering amplitude in the center-of-mass frame, where  $p_1 = (\sqrt{m^2 + \vec{p}^2}, \vec{p})$ ,  $p_2 = (\sqrt{m^2 + \vec{p}^2}, -\vec{p})$ ,  $p_3 = (\sqrt{m^2 + \vec{p}'^2}, \vec{p}')$ ,  $p_4 = (\sqrt{m^2 + \vec{p}'^2}, -\vec{p}')$ . Let  $P = p_1 + p_2 = (2p_0, \vec{0})$  denote the total four-momentum of the scattered nucleons where  $p_0 = \sqrt{\vec{p}^2 + m^2}$ .

The leading-order effective potential consists of the contact interaction part and the one-pion-exchange diagram. The leading-order contact interaction potential reads

$$\begin{aligned} iV_{0,C;\lambda\sigma,\mu\nu}^{i_3 i_4, i_1 i_2} = & 2i \sum_{a=0}^3 \left\{ C_S^a (\delta_{\lambda\nu} \delta_{\sigma\mu} \tau_{i_3 i_2}^a \tau_{i_4 i_1}^a - \delta_{\lambda\mu} \delta_{\sigma\nu} \tau_{i_3 i_1}^a \tau_{i_4 i_2}^a) \right. \\ & + C_A^a (\gamma_{5;\lambda\nu} \gamma_{5;\sigma\mu} \tau_{i_3 i_2}^a \tau_{i_4 i_1}^a - \gamma_{5;\lambda\mu} \gamma_{5;\sigma\nu} \tau_{i_3 i_1}^a \tau_{i_4 i_2}^a) \\ & + C_V^a (\gamma_{\lambda\nu}^\alpha \gamma_{\alpha;\sigma\mu} \tau_{i_3 i_2}^a \tau_{i_4 i_1}^a - \gamma_{\lambda\mu}^\alpha \gamma_{\alpha;\sigma\nu} \tau_{i_3 i_1}^a \tau_{i_4 i_2}^a) \\ & + C_{AV}^a \left[ (\gamma_5 \gamma_\alpha)_{\lambda\nu} (\gamma_5 \gamma^\alpha)_{\sigma\mu} \tau_{i_3 i_2}^a \tau_{i_4 i_1}^a - (\gamma_5 \gamma_\alpha)_{\lambda\mu} (\gamma_5 \gamma^\alpha)_{\sigma\nu} \tau_{i_3 i_1}^a \tau_{i_4 i_2}^a \right] \\ & \left. + C_T^a \left[ (\sigma^{\alpha\beta})_{\lambda\nu} (\sigma_{\alpha\beta})_{\sigma\mu} \tau_{i_3 i_2}^a \tau_{i_4 i_1}^a - (\sigma^{\alpha\beta})_{\lambda\mu} (\sigma_{\alpha\beta})_{\sigma\nu} \tau_{i_3 i_1}^a \tau_{i_4 i_2}^a \right] \right\}. \quad (34) \end{aligned}$$

The projected potential is

$$i\tilde{V}_{0,C;ls,mn}^{i_3i_4,i_1i_2} = 2i \sum_{a=0}^3 \left\{ (C_S^a + C_V^a) (\delta_{ln}\delta_{sm}\tau_{i_3i_2}^a\tau_{i_4i_1}^a - \delta_{lm}\delta_{sn}\tau_{i_3i_1}^a\tau_{i_4i_2}^a) \right. \\ \left. - (C_{AV}^a + 2C_T^a) \sum_{b=1}^3 [\sigma_{ln}^b\sigma_{sm}^b\tau_{i_3i_2}^a\tau_{i_4i_1}^a - \sigma_{lm}^b\sigma_{sn}^b\tau_{i_3i_1}^a\tau_{i_4i_2}^a] \right\}, \quad (35)$$

with  $l, s, m, n = 1, 2$ . Note that the pseudoscalar piece of the leading-order NN Lagrangian starts contributing at higher orders.

The covariant form of the one-pion-exchange potential reads

$$iV_{0,OPE;\lambda\sigma,\mu\nu}^{i_3i_4,i_1i_2}(p_3, p_4, p_1, p_2) = - \sum_{a=1}^3 i \frac{g_A^2}{4F^2} \frac{[\gamma_5(\not{p}_1 - \not{p}_3)]_{\lambda\mu} [\gamma_5(\not{p}_1 - \not{p}_3)]_{\sigma\nu}}{(p_1 - p_3)^2 - M^2} \tau_{i_3i_1}^a \tau_{i_4i_2}^a \\ + \sum_{a=1}^3 i \frac{g_A^2}{4F^2} \frac{[\gamma_5(\not{p}_1 - \not{p}_4)]_{\sigma\mu} [\gamma_5(\not{p}_1 - \not{p}_4)]_{\lambda\nu}}{(p_1 - p_4)^2 - M^2} \tau_{i_3i_2}^a \tau_{i_4i_1}^a. \quad (36)$$

In order to simplify some technical aspects of solving the leading-order equation we separate the pion denominator as

$$\frac{1}{k^2 - M^2} = -\frac{1}{\vec{k}^2 + M^2} + \left( \frac{1}{k^2 - M^2} + \frac{1}{\vec{k}^2 + M^2} \right), \quad (37)$$

and take the first term on the right-hand side of Eq. (37) as the leading-order contribution in the effective potential.<sup>1</sup> The second term is of higher order and is taken into account perturbatively. The resulting OPE potential reads

$$iV_{0,\pi;\lambda\sigma,\mu\nu}^{i_3i_4,i_1i_2}(\vec{p}', \vec{p}) = \sum_{a=1}^3 i \frac{g_A^2}{4F^2} \frac{[\gamma_5(\not{p}_1 - \not{p}_3)]_{\lambda\mu} [\gamma_5(\not{p}_1 - \not{p}_3)]_{\sigma\nu}}{(\vec{p}_1 - \vec{p}_3)^2 + M^2} \tau_{i_3i_1}^a \tau_{i_4i_2}^a \\ - \sum_{a=1}^3 i \frac{g_A^2}{4F^2} \frac{[\gamma_5(\not{p}_1 - \not{p}_4)]_{\sigma\mu} [\gamma_5(\not{p}_1 - \not{p}_4)]_{\lambda\nu}}{(\vec{p}_1 - \vec{p}_4)^2 + M^2} \tau_{i_3i_2}^a \tau_{i_4i_1}^a. \quad (38)$$

The projection reduces Eq. (38) to the following form,

$$i\tilde{V}_{0\pi;ls,mn}^{i_3i_4,i_1i_2}(\vec{p}', \vec{p}) = \sum_{a=1}^3 i \frac{g_A^2}{4F^2} \frac{[\vec{\sigma} \cdot (\vec{p} - \vec{p}')]_{lm} [\vec{\sigma} \cdot (\vec{p} - \vec{p}')]_{sn}}{(\vec{p} - \vec{p}')^2 + M^2} \tau_{i_3i_1}^a \tau_{i_4i_2}^a \\ - \sum_{a=1}^3 i \frac{g_A^2}{4F^2} \frac{[\vec{\sigma} \cdot (\vec{p} + \vec{p}')]_{sm} [\vec{\sigma} \cdot (\vec{p} + \vec{p}')]_{ln}}{(\vec{p} + \vec{p}')^2 + M^2} \tau_{i_3i_2}^a \tau_{i_4i_1}^a. \quad (39)$$

The leading-order scattering amplitude satisfies the equation

$$\tilde{T}_{0;ls,mn}^{i_3i_4,i_1i_2}(p_3, p_4, p_1, p_2) = \tilde{V}_{0;ls,mn}^{i_3i_4,i_1i_2}(\vec{p}', \vec{p}) - i \int \frac{d^4k}{(2\pi)^4} \tilde{V}_{0;ls,ab}^{i_3i_4,ij}(\vec{p}', -\vec{k})$$

---

<sup>1</sup> Without this simplification the Bethe-Salpeter equation has to be solved, which is technically more involved.




FIG. 1: Two examples of iterations of the LO equation.

$$\begin{aligned}
& \times \frac{m^2 F^2(L^2, \vec{k}^2)}{(\vec{k}^2 + m^2) (p_0 - k_0 - \sqrt{\vec{k}^2 + m^2} + i\epsilon) (p_0 + k_0 - \sqrt{\vec{k}^2 + m^2} + i\epsilon)} \\
& \times \tilde{T}_{0;ab,mn}^{ij,i_1i_2}(P/2 - k, P/2 + k, p_1, p_2), \quad (40)
\end{aligned}$$

where

$$\tilde{V}_0 = \tilde{V}_{0,C} + \tilde{V}_{0,\pi}. \quad (41)$$

As the leading-order potential does not depend on  $k_0$ , the integration over this variable can be performed easily by iterating Eq. (40), performing the  $k_0$  integration in each term and summing up the resulting terms. Doing so we obtain

$$\begin{aligned}
\tilde{T}_{0;ls,mn}^{i_3i_4,i_1i_2}(\vec{p}', \vec{p}) &= \tilde{V}_{0;ls,mn}^{i_3i_4,i_1i_2}(\vec{p}', \vec{p}) - \frac{m^2}{2} \int \frac{d^3k}{(2\pi)^3} \tilde{V}_{0;ls,ab}^{i_3i_4,ij}(\vec{p}', -\vec{k}) \\
&\times \frac{F^2(L^2, \vec{k}^2)}{(\vec{k}^2 + m^2) (p_0 - \sqrt{\vec{k}^2 + m^2} + i\epsilon)} \tilde{T}_{0;ab,mn}^{ij,i_1i_2}(-\vec{k}, \vec{p}). \quad (42)
\end{aligned}$$

Next we split the two-nucleon propagator into two parts,

$$\begin{aligned}
\frac{1}{(\vec{k}^2 + m^2) (p_0 - \sqrt{\vec{k}^2 + m^2} + i\epsilon)} &= \frac{2m}{(\vec{k}^2 + m^2) (\vec{p}^2 - \vec{k}^2 + i\epsilon)} \\
&+ \frac{(p_0 + \sqrt{\vec{k}^2 + m^2}) - 2m}{(\vec{k}^2 + m^2) (\vec{p}^2 - \vec{k}^2 + i\epsilon)}, \quad (43)
\end{aligned}$$

and take the first term as the leading order propagator while including the second part in perturbative corrections. The resulting leading-order equation reads

$$\begin{aligned}
\tilde{T}_{0;ls,mn}^{i_3i_4,i_1i_2}(\vec{p}', \vec{p}) &= \tilde{V}_{0;ls,mn}^{i_3i_4,i_1i_2}(\vec{p}', \vec{p}) - m^3 \int \frac{d^3k}{(2\pi)^3} \tilde{V}_{0;ls,ab}^{i_3i_4,ij}(\vec{p}', -\vec{k}) \\
&\times \frac{F^2(L^2, \vec{k}^2)}{(\vec{k}^2 + m^2) (\vec{p}^2 - \vec{k}^2 + i\epsilon)} \tilde{T}_{0;ab,mn}^{ij,i_1i_2}(-\vec{k}, \vec{p}). \quad (44)
\end{aligned}$$

Let us investigate the "consistency problem" which manifests itself in the sensitivity of the scattering amplitude to the choice of the parameter  $L$ . For that purpose we vary  $L$  from 750 MeV to infinity. Divergences which occur in the limit  $L \rightarrow \infty$  will be referred to as UV

divergences. Simple UV counting shows that iterations of Eq. (44) lead to finite diagrams for  $L \rightarrow \infty$ . For example, the diagrams in Fig. 1 are finite. This differs from the standard non-relativistic formulation, in which Fig. 1 (a) contains an overall logarithmic divergence  $\sim M^2 m^2$  and the diagram in Fig. 1 (b) contains a logarithmic divergence  $\sim (Qm)^6$ . The divergences in these diagrams can be absorbed in contact interaction terms of the second and sixth order (and higher-order iterations require higher-order interaction terms, up to infinity) [9].

Equation (44) can be rewritten in a more familiar form. To that end we multiply Eq. (44) with

$$\frac{m^2 F(L^2, \vec{p}'^2) F(L^2, \vec{p}^2)}{[(m^2 + \vec{p}'^2)(m^2 + \vec{p}^2)]^{1/2}}$$

and obtain

$$\begin{aligned} \mathcal{T}_{0;ls,mn}^{i_3 i_4, i_1 i_2}(\vec{p}', \vec{p}) &= \mathcal{V}_{0;ls,mn}^{i_3 i_4, i_1 i_2}(\vec{p}', \vec{p}) \\ &- m \int \frac{d^3 k}{(2\pi)^3} \mathcal{V}_{0;ls,ab}^{i_3 i_4, ij}(\vec{p}', -\vec{k}) \frac{1}{\vec{p}^2 - \vec{k}^2 + i\epsilon} \mathcal{T}_{0;ab,mn}^{ij, i_1 i_2}(-\vec{k}, \vec{p}), \end{aligned} \quad (45)$$

where

$$\begin{aligned} \mathcal{T}_0(\vec{p}', \vec{p}) &= \frac{\tilde{T}_0(\vec{p}', \vec{p}) F(L^2, \vec{p}'^2) F(L^2, \vec{p}^2) m^2}{[(m^2 + \vec{p}'^2)(m^2 + \vec{p}^2)]^{1/2}} \\ \mathcal{V}_0(\vec{p}', \vec{p}) &= \frac{\tilde{V}_0(\vec{p}', \vec{p}) F(L^2, \vec{p}'^2) F(L^2, \vec{p}^2) m^2}{[(m^2 + \vec{p}'^2)(m^2 + \vec{p}^2)]^{1/2}}. \end{aligned} \quad (46)$$

Note that the potential  $\mathcal{V}_0(\vec{p}, \vec{q})$  has a milder ultraviolet behavior than the standard potential  $\tilde{V}_0(\vec{p}', \vec{p})$  due to the denominator in Eq. (46). In the limit  $L \rightarrow \infty$  the factors  $F(L^2, \vec{p}^2)$  and  $F(L^2, \vec{p}'^2)$  approach 1. The OPE part of the effective potential  $\mathcal{V}_0(\vec{p}', \vec{p})$  of Eq. (46) then behaves like  $\sim 1/p^3$  for large  $p$  in the  $^1S_0$  wave (after absorbing the local part in the redefinition of the contact interaction term) and generates no divergences (perturbatively as well as non-perturbatively). This is in contrast to the OPE part of  $\tilde{V}_0(\vec{p}', \vec{p})$  which goes as  $\sim 1/p^2$ . The most singular part of the effective OPE potential in the  $^3S_1$  wave requires a single counter-term, which is present in the contact interaction part of our LO potential. For higher partial waves the singular behavior of the potential is screened by the angular momentum barrier and therefore no counter-terms are required. Analyzing the convergence criterium for Neumann series we have checked that the integral equations for higher partial waves have unique solutions, i.e. the angular momentum barrier provides a repulsion strong enough for our problem. The numerical analysis confirms the above conclusions. Some of these phase shifts calculated for different values of the parameter  $L$  are shown in Figs. 2, 3 and 4. Note that different choices of the parameter  $L$  correspond to different renormalization schemes. By choosing the renormalization scheme properly one can improve the convergence of perturbative series for physical quantities. Our figures suggest that the best description of the phase shifts at leading order is obtained for  $L \sim 750$  MeV. This indicates that the higher-order corrections are smaller for this choice of the renormalization condition.

We conclude that all counter-terms required for the absorbition of the strong  $L$  dependence in the LO equation (42), perturbatively as well as non-perturbatively, are present in the LO potential. Therefore, Eq. (42) is free of the "problems of consistency of Weinberg's power




FIG. 2:  $^1S_0$  partial wave  $np$  phase shifts for different values of  $L$  parameter compared with the data points from the Nijmegen PWA [31]. The curves are shown for  $L = 750, 1000, 2000, 6000, 20000, 30000, 100000$  (MeV), the solid line corresponding to the largest value.




FIG. 3:  $^3S_1$  partial wave  $np$  phase shifts for different values of  $L$  parameter compared with the data points from the Nijmegen PWA [31]. The curves are shown for  $L = 750, 1000, 2000, 6000, 10000, 20000, 50000$  (MeV), the solid line corresponding to the largest value.

counting” encountered in solving the corresponding Lippmann-Schwinger equation [6, 7, 9].<sup>2</sup>

Note that Eq. (42) in comparison with the LS equation contains the re-summation of an infinite number of higher-order terms. In the authors’ opinion both approaches are equivalent and self-consistent, provided that one includes compensating terms in the effective Lagrangian and chooses the cutoff parameter appropriately in the LS approach. Therefore we expect the numerical results for physical quantities in our approach to be comparable to that of the cutoff LS equation approach [5].

---

<sup>2</sup> The Lippmann-Schwinger equation corresponds to the leading-order term in the  $1/m$  expansion of Eq. (42).




FIG. 4:  ${}^3P_0$  partial wave  $np$  phase shifts for different values of  $L$  parameter compared with the data points from the Nijmegen PWA [31]. The curves are shown for  $L = 750, 1000, 2000, 6000, 20000, 30000, 100000, 500000$  (MeV), the solid line corresponding to the largest value.

### B. Higher-order Corrections

Corrections to the leading-order amplitude are calculated perturbatively and to any finite order can be expressed in a closed form as a sum of a finite number of terms analogous to Eq. (33). These expressions can be interpreted as sums of a finite number of Feynmann diagrams where the leading-order amplitude  $T_0$  is also interpreted as an NN vertex. The straightforward analysis of Eqs. (44)-(45) shows that (for  $L \rightarrow \infty$ )  $T_0(\vec{p}, \vec{p}')$  cannot diverge stronger than  $\sim \ln p$  ( $\sim \ln p'$ ) for  $p \rightarrow \infty$  ( $p' \rightarrow \infty$ ). This UV behavior of  $T_0$  guarantees that all divergences generated in the limit  $L \rightarrow \infty$  in Feynman diagrams contributing to corrections at any finite order can be canceled by counter-term contributions also present at the given order. As a result the new semi-relativistic approach suggested in this work is free of "consistency problems" inherent in the standard non-relativistic approach.

## V. SUMMARY

We consider a new approach to the nucleon-nucleon scattering problem in baryon chiral perturbation theory. Starting with a Lorentz-invariant form of the effective Lagrangian with higher derivative terms we work out a new framework where the nucleon-nucleon scattering amplitude is calculated order-by-order in small parameters like the pion mass and nucleon three-momenta. We develop a systematic power counting for the effective potential which is defined as the sum of all two-nucleon irreducible diagrams. The leading-order amplitude is obtained by solving an equation with the LO potential consisting of the contact interaction plus OPE. Higher-order corrections to the potential as well as to the two-nucleon propagator are taken into account perturbatively. The parameter  $L$ , introduced through terms with higher derivatives, parameterizes the renormalization scheme dependence of the scattering amplitude present in our calculations. Due to the UV behavior of our semi-relativistic two-nucleon propagator, analogously to the KSW approach, the strong  $L$ -dependence of physical quantities to any finite order is absorbed in the redefinition of a finite number of parameters of the effective potential at given order. On the other hand, analogously to Weinberg's

power counting, the OPE potential is of leading order and is treated non-perturbatively. Our new approach preserves all underlying symmetries to the given order of accuracy and the self-consistency of the power counting is explicit in this new formulation.

## Acknowledgments

D.D., J.G. and M.R.S. acknowledge the support of the Deutsche Forschungsgemeinschaft (SFB 443).

- 
- [1] S. Weinberg, Phys. Lett. B **251**, 288 (1990).
  - [2] S. Weinberg, Nucl. Phys. **B363**, 3 (1991).
  - [3] C. Ordonez and U. van Kolck, Phys. Lett. B **291**, 459 (1992).
  - [4] P. F. Bedaque and U. van Kolck, Ann. Rev. Nucl. Part. Sci. **52**, 339 (2002).
  - [5] E. Epelbaum, Prog. Part. Nucl. Phys. **57**, 654 (2006).
  - [6] D. B. Kaplan, M. J. Savage, and M. B. Wise, Nucl. Phys. **B478**, 629 (1996).
  - [7] D. B. Kaplan, M. J. Savage, and M. B. Wise, Phys. Lett. B **424**, 390 (1998).
  - [8] D. B. Kaplan, M. J. Savage, and M. B. Wise, Nucl. Phys. **B534**, 329 (1998).
  - [9] M. J. Savage, arXiv:nucl-th/9804034.
  - [10] J. Gegelia, arXiv:nucl-th/9806028.
  - [11] J. Gegelia, Phys. Lett. B **463**, 133 (1999).
  - [12] T. D. Cohen and J. M. Hansen, Phys. Rev. C **59**, 3047 (1999).
  - [13] S. Fleming, T. Mehen, and I. W. Stewart, Nucl. Phys. **A677**, 313 (2000).
  - [14] S. R. Beane, P. F. Bedaque, M. J. Savage, and U. van Kolck, Nucl. Phys. **A700**, 377 (2002).
  - [15] A. Nogga, R. G. E. Timmermans, and U. van Kolck, Phys. Rev. C **72**, 054006 (2005).
  - [16] E. Epelbaum, W. Glöckle, and U. -G. Meißner, Nucl. Phys. **A747**, 362 (2005).
  - [17] G. P. Lepage, arXiv:nucl-th/9706029.
  - [18] J. Gegelia, Phys. Lett. B **429**, 227 (1998).
  - [19] J. Gegelia, J. Phys. G **25**, 1681 (1999).
  - [20] T. S. Park, K. Kubodera, D. P. Min, and M. Rho, Nucl. Phys. **A646**, 83 (1999).
  - [21] G. P. Lepage, *Conference summary*, Prepared for INT Workshop on Nuclear Physics with Effective Field Theory, Seattle, Washington, 25-26 Feb 1999.
  - [22] J. Gegelia and S. Scherer, Int. J. Mod. Phys. A **21**, 1079 (2006).
  - [23] E. Epelbaum and U. -G. Meißner, arXiv:nucl-th/0609037.
  - [24] D. Djukanovic, M. R. Schindler, J. Gegelia, and S. Scherer, Phys. Rev. D **72**, 045002 (2005).
  - [25] J. Gasser and H. Leutwyler, Ann. Phys. (N.Y.) **158**, 142 (1984).
  - [26] J. Gasser, M. E. Sainio, and A. Švarc, Nucl. Phys. **B307**, 779 (1988).
  - [27] S. Scherer and H. W. Fearing, Phys. Rev. D **52**, 6445 (1995).
  - [28] A. A. Bel'kov, A. V. Lanyov, A. Schaale, and S. Scherer, Acta Phys. Slov. **45**, 121 (1995).
  - [29] N. Fettes, U.-G. Meißner, M. Mojžiš, and S. Steininger, Ann. Phys. (N.Y.) **283**, 273 (2000); **288**, 249 (2001).
  - [30] G. Ecker, Prog. Part. Nucl. Phys. **35**, 1 (1995).
  - [31] V. G. J. Stoks, R. A. M. Klomp, M. C. M. Rentmeester and J. J. de Swart, Phys. Rev. C **48**, 792 (1993), <http://nn-online.org>.